STD PREVENTION AND CONTROL SERVICES, SAN FRANCISCO DEPARTMENT OF PUBLIC HEALTH STD AND HIV SCREENING GUIDELINES

These evidence-based recommendations provide guidance for chlamydia, gonorrhea, syphilis and HIV screening in persons without symptoms or a need for diagnostic testing.¹

WOMEN

	Chlamydia and Gonorrhea	Syphilis	HIV
25 years and younger	Test every 12 months	Not routinely recommended ²	At least one lifetime test ²
Older than 25 years	Not routinely recommended ²		At least one lifetime test ²
Pregnant ³	Test in 1 st trimester, repeat in 3 rd trimester if at increased risk ²	Test in 1 st trimester, repeat in 3 rd trimester if at increased risk ²	First prenatal visit, repeat in 3 rd trimester if at increased risk ²
HIV+ Women should be screened annually for trichomoniasis (trich)			

MEN WHO HAVE SEX WITH WOMEN

	Chlamydia and Gonorrhea	Syphilis	HIV
Any age, any site	Not routinely recommended ²		At least one lifetime test ²

MEN WHO HAVE SEX WITH MEN (MSM)

men with the sex with men (mon)					
	Chlamydia and Gonorrhea	Syphilis	HIV		
Blood		Every 3 months			
Rectal & Pharyngeal ⁴	Every 3 months				
Urine⁵	Every 3 months				

TRANS WOMEN AND TRANS MEN WHO HAVE SEX WITH MEN

THE WAY OF THE PROPERTY OF THE				
	Chlamydia and Gonorrhea	Syphilis	HIV	
Blood		Every 3 months		
Rectal & Pharyngeal ⁴	Every 3 months			
Urine⁵/Vaginal Swab	Every 3 months			

¹These are general population recommendations. SFDPH may make separate recommendations for specific groups that correlate with higher risk.

Terminology Defined: Transgender (Trans): A person whose gender identity differs from the sex that was assigned at birth. A trans man (or trans male) is someone with a male gender identity who was assigned female sex at birth; a trans woman (or trans female) is someone with a female gender identity who was assigned male sex at birth.

For more info: Visit www.sfcityclinic.org or call 415-487-5595

²Consider screening if patient has any of the following risk factors: sex with a man who has sex with men, history of STD in the past year, methamphetamine use, unstable housing or homelessness, sex work, intimate partner violence, or incarceration.

³Regardless of intentions to carry to term.

⁴Prioritize these extra-genital sites, as rectal and pharyngeal infections are almost always asymptomatic.

⁵If cost permits or extra-genital testing not available.