

Patient Delivered Partner Therapy for Gonorrhea and Chlamydia (Azithromycin 1 g and Cefixime 800 mg)

PLEASE READ THIS VERY CAREFULLY

Why do I need to take this medicine?

Your sex partner has recently been treated for Gonorrhea and Chlamydia. These are curable infections you can get from having sex with a person who already has them. Many people with these infections do not know they have them because they feel okay and do not have any symptoms. However, if you do not take medicine to cure it, you can get very sick.

You could have Gonorrhea and/or Chlamydia. It is important that you get treated. We want to be sure that you get the medicine you need as soon as possible. ***The best treatment for Chlamydia and Gonorrhea includes an antibiotic pill plus an antibiotic injection. You can only receive the injection at City Clinic or a doctor's office.*** If you are not able to go to City Clinic or your doctor within 1 week, you should take the medicine enclosed.

Is the medicine safe?

The medicine is very safe. However, you should talk to your doctor or come to City Clinic if you have any of the problems listed below. DO NOT take the medicine if:

- You are female and having lower belly pain, pain with sex, vomiting or fever
- You are male and having pain or swelling in the testicle (balls) or fever
- You ever had a bad reaction, rash or allergy to Azithromycin (Zithromax), Erythromycin, or Clarithromycin (Biaxin)
- You ever had a bad reaction, rash or allergy to Cefixime (Suprax), Cefpodoxime (Vantin) or any cephalosporin (any antibiotic that starts with cef- or cep-) or a life threatening reaction to Penicillin
- You have a serious long-term illness such as kidney, heart or liver disease
- You are taking prescribed medicine, especially for heart disease or serious mental illness

How do I take the medicine?

- You will receive 2 pink or blue 500 mg Azithromycin pills (for Chlamydia and Gonorrhea) and 2 white 400 mg Cefixime pills (for Gonorrhea).
- Take the Azithromycin and Cefixime pills at the same time with food and a full glass of water.
- This medicine may cause an upset stomach, diarrhea, nausea, or vomiting. This should not last long. If these symptoms persist or you develop a rash or fever, contact your doctor or City Clinic.
- Do not share or give this medicine to anyone else.
- Do not have sex for the next 7 days because it takes 7 days for the medicine to cure these infections. If you have unprotected sex during the 7 days after taking the medicine, you could still pass the infection to your sex partners.
- We recommend re-testing in 3 months to make sure that you have not been re-infected.

What can I do if I still have questions?

If you have any questions about the medicine, please call (415) 487-5595 during clinic hours. All calls are confidential. You can also come into City Clinic for a visit including exam, testing and medicine. For more information and directions, visit our website.

City Clinic
356 7th Street
San Francisco, CA 94103
(415) 487-5500
www.sfcityclinic.org

Monday	8 am – 4pm
Tuesday	1 pm – 6 pm
Wednesday	8 am – 4 pm
Thursday	1 pm – 4 pm
Friday	8 am – 4 pm

